75. Number of young-of-year striped bass captured by semi-balloon trawl in western Albemarle Sound, NC, by station, July-October 1992 218

76. JAI catch matrix for seven stations in western Albemarle Sound, NC, 1992 219

77. Number of young-of-year striped bass captured by semi-balloon trawl in western Albemarle Sound, NC, by station, July-October, 1993 220

78. JAI catch matrix for seven stations in western Albemarle Sound, NC, 1993 220

79. Description of trawl and beach seine sampling stations in Albemarle Sound used by the N.C. Division of Marine Fisheries 228

80. Comparisons of striped bass spawning activity in the Roanoke River documented by field observations of the egg study, and by backcalculating spawn dates of otoliths from juveniles collected by the Hassler trawl survey (JAI), the central and eastern trawl survey (EST), the alosid beach seine survey (ALO), and the exploratory beach seine survey (EXP) 234

81. Percent occurrence of prey taxa in striped bass stomachs from western and eastern Albemarle Sound, NC, in 1991 242

82. Density (number/100 m^3) of striped bass larvae in the lower Roanoke River (Stations 1-12), North Carolina, 1990-1991 245

83. Density (number/100 m^3) of striped bass larvae in Batchelor Bay (Stations 13-16) and western Albemarle Sound (Stations 17-32), North Carolina, 1990-1991 246

84. Relative contribution (% by enumeration) of prey consumed by larval striped bass in the lower Roanoke River (Stations 1-12), Batchelor Bay (Stations 13-16), and western Albemarle Sound (Stations 17-32), North Carolina, 1984-1991 248

85. Date at which feeding by striped bass larvae was first observed in the lower Roanoke River (Stations 1-12), Batchelor Bay (Stations 13-16), and western Albemarle Sound, North Carolina, 1984-1991 249

86. Mean catch per trawl of 10 finfish species other than striped bass collected in western Albemarle Sound by NC Division of Marine Fisheries personnel, 1982-1987 and 1988-1993 256

87. Catch per unit effort (number of fish/trawl) of selected finfish species collected in western Albemarle Sound by N.C. Division of Marine Fisheries personnel, 1982-1993 257

88. Relative contribution (% using density) of each taxonomic group to the spring zooplankton community of the lower Roanoke River (Stations 1-12), Batchelor Bay (Stations 13-16), and western Albemarle Sound (Stations 17-32), North Carolina, 1990-1991 269

89. Results of linear regressions to determine the relationship of body width to body length of seven larval fish taxa 272